

## ON HIGHER DIMENSIONAL ORCHARD VISIBILITY PROBLEM

SHENGNING ZHANG

ABSTRACT. In this article, we study Pólya's orchard visibility problem in arbitrary dimension  $d$ : suppose at every integral point in  $\mathbb{R}^d$ , centered a small  $d$ -dimensional ball with radius  $r$  (which is considered as a tree at the integral point), given a  $d$ -dimensional ball centered at the origin  $O$  with radius  $R$  (which is considered as the orchard), it asks for the smallest  $r$  such that every ray starting from  $O$  will hit some tree in the orchard. We give both upper and lower bounds of the minimal value of  $r$ , say  $\rho$  in terms of  $R$ , moreover, we prove that as  $R \rightarrow \infty$ ,  $\rho = O(R^{-\frac{1}{d-1}})$ .

## 1. INTRODUCTION

Let  $\Lambda$  be the set of lattice points  $\mathbb{Z}^d \setminus O$  in  $\mathbb{R}^d$ , where  $O$  is the origin. Let  $B(O, R)$  be the closed ball in  $\mathbb{R}^d$  centered at  $O$  with radius  $R > 1$ . Centering at every integral point  $P \in B(O, R)$ , is a small closed ball  $B(P, r)$  with given small radius  $r > 0$ . The original Pólya's orchard visibility problem considers the case  $d = 2$ , when the disc  $B(O, R)$  is thought as a round orchard and every  $B(P, r)$  a tree at  $P$ , it asks for the smallest  $r$ , which we denote by  $\rho$ , so that one standing at the center  $O$  *cannot* see through the orchard, that is, for any ray  $l$  starting from  $O$ ,  $l \cap B(P, r) \neq \emptyset$  for some  $P$ .

In [1], it proved that

$$(1.1) \quad \frac{1}{\sqrt{R^2+1}} < \rho < \frac{1}{R}.$$

Indeed, in an earlier paper [2], Thomas Tracy Allen had proved that

$$(1.2) \quad \rho = \frac{1}{R}.$$

In this paper, we'd like to study the general Pólya's orchard problem in arbitrary dimension  $d$  and prove similar bounds as in (1.1). Our strategy follows [3], where, however, only deals with the 2 and 3 dimensional cases.

## 2. LOWER BOUNDS

Consider in  $\mathbb{R}^d$  the  $d$ -dimensional cuboid  $C$  with diagonal vertices  $O$  and  $D := (1, 1, \dots, 1, [R] + 1)$ , where  $[R]$  is the floor function of  $R$ . Then

$$C \cap \mathbb{Z}^d = \{(x_1, \dots, x_d) \in \mathbb{Z}^d \mid x_i \in [0, 1], \forall i = 1, \dots, d-1; x_d \in [0, [R] + 1]\}.$$

Apparently,  $D$  is not in  $B(O, R)$ . The segment  $OD$  is of the length  $\sqrt{(d-1) + R^2}$ , and any  $P \in C \cap \mathbb{Z}^d$  has the distance squared  $dist(P, OD)^2$  to  $OD$

$$(2.1) \quad \frac{(d-1 + ([R] + 1)^2)(x_1^2 + \dots + x_d^2) - (x_1 + \dots + x_{d-1} + ([R] + 1)x_d)^2}{d-1 + ([R] + 1)^2}$$

This lead us to our first result, which is a direct generalization to the first inequality of (1.1).

PROPOSITION 1. *notations as above*

$$(2.2) \quad \frac{\sqrt{d-1}}{\sqrt{d-1 + ([R] + 1)^2}} < \rho.$$

PROOF: Consider the formula (2.1), apparently that among all integral points in  $C$  other than  $O$  and  $D$ ,  $P_0 = \{0, \dots, 0, 1\}$  minimize the expression, when

$$dist(P_0, OD)^2 = \frac{d-1}{d-1 + ([R] + 1)^2}.$$

(see the figure below)




FIGURE 1

So if the tree radius  $r$  can block the orchard, it must bigger than  $\frac{\sqrt{d-1}}{\sqrt{d-1 + ([R] + 1)^2}}$ . This completes the proof.  $\square$

The proposition tells us that  $\rho$  grows faster than the rate of  $R^{-1}$  as  $R$  goes to infinity, however, it is not the exact rate of growth of  $\rho$ , so we want a better

lower bound of  $\rho$  in terms of  $R$ . Indeed, the proof of the proposition tells us that, to obtain such a lower bound, we have to consider a “finer” solid containing the ray than the coboid  $C$  above. To use such a solid in higher dimension, we have to use the volume formula of a lattice polyhedron in higher dimension developed by Macdonald in [4], which is the generalization of Pick’s Theorem used in [3, Theorem 2.2]. Now we summarize below.

Let  $\mathbb{Z}^d \subset \mathbb{R}^d$  be the standard integral lattice,  $X$  a  $d$ -dimensional polyhedra in  $\mathbb{R}^d$  whose vertices are all in  $\mathbb{Z}^d$ . Let  $\partial X$  be the boundary of  $X$ , which can be viewed as a  $d - 1$ - simplicial complex. For any integer  $n > 0$ , write

$$L(n, X) = |X \cap \frac{1}{n}\mathbb{Z}^d|,$$

and

$$M(n, X) = L(n, X) - \frac{1}{2}L(n, \partial X),$$

then, we have the volume of  $X$  can be computed by:

PROPOSITION 2 (Macdonald’s Theorem). *The volume of the polyhedra  $Vol(X)$  equals*

$$\frac{2}{(d-1)d!} \left\{ M(d-1, X) - \binom{d-1}{1} M(d-2, X) + \binom{d-1}{2} M(d-3, X) - \dots + (-1)^{d-1} M(0, X) \right\},$$

where  $M(0, X) = 1$  if  $d$  is even,  $M(0, X) = 0$  if  $d$  is odd.

Now we give us first theorem

THEOREM 1. *There is a constant  $c > 0$  such that*

$$(2.3) \quad ([R] + 1)\rho^{d-1} > c.$$

*Remark 1.* The constant  $c$  is given by the volume of a polyhedra, which can be computed using Macdonald’s Theorem above. The key is to construct a proper polyhedra, which will be clear in the proof of the theorem.

LEMMA 1. *Point  $Q \in \mathbb{Z}^d \cap B(O, R)$ , if for any  $P \in \mathbb{Z}^d \cap B(O, R)$ ,  $OB \cap B(P, r) = \emptyset$ , then the coordinates of  $Q$  are coprime, that is, if  $Q = (a_1, \dots, a_d)$  then  $\gcd(a_1, \dots, a_d) = 1$ .*

The lemma comes from an easy observation. Suppose  $\gcd(a_1, \dots, a_d) = d > 1$ , then  $P_1 = \frac{1}{d}(a_1, \dots, a_d) \in \mathbb{Z}^d \cap B(O, R)$  and obviously  $OB \cap B(P_1, r) \neq \emptyset$ .  $\square$

LEMMA 2. *Let  $l$  be any ray starting from  $O$ , if point  $P \in \mathbb{Z}^d \cap B(O, R)$ ,  $P \notin l$  such that  $\text{dist}(P, l)$  is minimal, then the coordinates of  $P$  are coprime.*

Suppose the coordinates of  $P$  are coprime with greatest common divisor  $d > 1$ , then  $\text{dist}(\frac{1}{d}P, l) < \text{dist}(P, l)$ . Contradiction.  $\square$

To carry out our argument in high dimension, we have to generalize the result to Lemma 2 from a ray  $l$  to a family of geometric objects which we called diamonds with a diagonal, and is defined as follow:

DEFINITION 1. In  $\mathbb{R}^d$ , for any positive integer  $n \leq d$ , a  $n$ -dimensional diamond  $\mathfrak{D}$  with a diagonal  $I$  is defined as follow:

- (i) A 1-dimensional diamond  $\mathfrak{D}$  is nothing but a segment start from the origin  $O$  to a point  $P \neq O$  in  $\mathbb{R}^d$  and its diagonal  $I$  is itself;
- (ii) Suppose for any  $i \leq n$ , the  $i$ -dimensional diamonds with a diagonal are well-defined, then a  $n$ -dimensional diamond  $\mathfrak{D}_n$  with a diagonal  $I_n$  is defined base on some  $n$ -dimensional diamond  $\mathfrak{D}_{n-1}$  with a diagonal  $I_{n-1}$ : let  $V_{n-1}$  be the  $n-1$  vector space generated by vectors in  $\mathfrak{D}_{n-1}$ , and  $P_n$  a point in  $\mathbb{R}^d \setminus V_{n-1}$ , consider  $OI_{n-1}$  and  $OP_n$  as two vectors, then define  $Q_n$  be the end point of the vector  $OI_{n-1} - OP_n$ , and  $\mathfrak{D}_n$  is defined to be the convex hull of  $\mathfrak{D}_{n-1} \cup \{P_n, Q_n\}$ , its diagonal is  $I_n := I_{n-1}$ .




FIGURE 2. an example of 1,2 and 3-diamonds

LEMMA 3. Let  $\mathfrak{D}$  be a  $n$ -dimensional diamond with a diagonal  $I$  in  $\mathbb{R}^d$ ,  $n < d$ ,  $V$  be  $n$ -dimensional subspace in  $\mathbb{R}^d$  generated by  $\mathfrak{D}$ . Now if a point  $P \in \mathbb{Z}^d \cap B(O, R)$ ,  $P \notin V$  such that  $\text{dist}(P, \mathfrak{D})$  is minimal, then the coordinates of  $P$  are coprime.

Suppose  $A \in \mathfrak{D}$  is the point such that  $\text{dist}(P, \mathfrak{D}) = \text{dist}(P, A) = a$ . Consider the triangle  $\triangle OAP$ , since  $\mathfrak{D}$  is a convex hull by the definition, the segment  $OA \subset \mathfrak{D}$ . Now if the greatest common divisor of the coordinates of  $P$  is  $m > 1$ ,

consider the point  $Q = \frac{1}{m}P \in OP$ . Find a point  $Q' \in OA \subset \mathfrak{D}$  such that  $QQ' \parallel AP$ , then apparently that  $dist(Q, \mathfrak{D}) < dist(P, \mathfrak{D})$ . Contradiction!  $\square$

PROOF OF THE THEOREM: Consider the point  $D_1 := D$  given above, we view the segment  $OD$  as a vector from the origin  $O$  to  $D$  and denote it by  $\vec{l}$ . Among all integral points in  $B(O, R)$ , find  $P_2$  in the first quadrant (that is, all the points are of nonnegative coordinates) be the one of minimal distance to  $\vec{l}$ . Write the minimal distance  $\varepsilon_1$ . From the lemmas above, we know the coordinates of  $P_2$  are coprime. View the segment  $OP_2$  as a vector and denote it by  $\vec{v}_1$ , and define vector  $\vec{u}_1 := \vec{l} - \vec{v}_1$ , define the two dimensional diamond  $D_2$  be the parallelogram spanned by  $\vec{v}_1$  and  $\vec{u}_1$ . From the two lemmas above,  $D_2$  does not contain any integral points of  $\Lambda$  other than the 4 vertices. Denote the 2-dimensional plane spanned by  $\vec{v}_1$  and  $\vec{u}_1$  by  $V_2$ . Using our notion of diamond,  $D_2$  is a 2-dimensional diamond with a diagonal  $l$ .

Now among all integral points in  $B(O, R) \setminus V_2$ , find one  $P_3$  in the first quadrant of the minimal distance to  $V_2 \cap B(O, R)$ . Write the minimal distance  $\varepsilon_2$ . Consider the 2-dimensional diamond  $D_2$  with diagonal  $\vec{l}$  and the point  $P_3$ , by Definition 1, they together define a 3-dimensional diamond  $D_3$  with diagonal  $\vec{l}$ . By Lemma 3, all the coordinates of  $P_3$  are coprime,  $D_3$  contains no integral points other than the 6 vertices. Denote the 3-dimensional vector space generated by vectors in  $D_3$  by  $V_3$ .

Keep this process, for all integer  $i = 1, 2, \dots, d$ , we obtain  $i$ -dimensional diamond  $D_i$  with diagonal  $\vec{l}$ ,  $V_i = span D_i$ , integral points  $P_i$  in the first quadrant such that

- (a)  $dist(P_i, V_{i-1} \cap V_{i-1}) = \varepsilon_{i-1}$  is minimal among all integral points in  $B(O, R) \setminus V_{i-1}$ ;
- (b)  $D_i$  is the diamond constructed by  $D_{i-1}$  and  $P_i$ ;
- (c)  $D_i$  contains no integral points other than its vertices.

It is easy to see, from our construction, the volume of  $D_i$  is

$$(2.4) \quad Vol(D_i) = \frac{2^{i-1}}{i!} \varepsilon_1 \cdots \varepsilon_{i-1} ([R] + 1).$$

In particular, Write  $\mathfrak{D} := D - d$ , its volume is

$$(2.5) \quad Vol(\mathfrak{D}) = \frac{2^{d-1}}{d!} \varepsilon_1 \cdots \varepsilon_{d-1} ([R] + 1),$$

which can also be calculated by Macdonald's formula as in Proposition 2. On the other hand, By our construction of  $\mathfrak{D}$ , if the tree radius  $r$  is such that every ray starting from  $O$  and passing through one point in  $\mathfrak{D}$  will be blocked by some tree, then  $r > \varepsilon_i$  for any  $i$ . So we have

$$(2.6) \quad \frac{2^{d-1}}{d!} r^{d-1} ([R] + 1) > Vol(\mathfrak{D}).$$

Writing

$$(2.7) \quad c = \frac{d! \text{Vol}(\mathfrak{D})}{2^{d-1}},$$

we complete the proof.  $\square$

*Remark 2.* If  $d = 2$ ,  $\text{Vol}(\mathfrak{D}) = \text{Vol}(D_2) = 1$ , then the Theorem tells that  $([R] + 1)\rho > 1$ , which reproduces the result in [3, Proposition 2.4]. If  $d = 3$ ,  $\text{Vol}(\mathfrak{D}) = \text{Vol}(D_3) = \frac{2}{3} \frac{\sqrt{2}}{2} \frac{\sqrt{2}}{2} = \frac{1}{3}$ , then the theorem tells that

$$(2.8) \quad ([R] + 1)\rho^2 > \frac{1}{2},$$

which is better than the result in [3, Proposition 4.4].

### 3. UPPER BOUNDS

In this section we give an upper bound of  $\rho$  in terms of  $R$ . The key ingredient is again Minkowski's theorem as [3, Theorem 4.1], which we summarize below.

**PROPOSITION 3** (Minkowski's Theorem). *Let  $m$  be a positive integer and  $F \subset \mathbb{R}^d$  a domain satisfying*

- (a)  $F$  is symmetric with respect to  $O$ ;
- (b)  $F$  is convex;
- (c)  $\text{Vol}(F) \geq m2^d$ .

*Then  $F$  contains at least  $m$  pairs of points  $\pm A_i \in \mathbb{Z}^d \setminus O$ ,  $1 \leq i \leq m$ , which are distinct from each other.*

Now we state an upper bound of  $\rho$ . The idea is essentially same to [3, §4], where, however, only deals with the 3-dimensional case.

**THEOREM 2.** *There is a constant  $C > 0$ , such that*

$$(3.1) \quad R\rho^{d-1} < C.$$

**PROOF:** For any diameter  $AA'$  of the ball  $B(O, R)$ , let's consider the  $d - 1$ -dimensional hyperellipsoid  $E \subset \mathbb{R}^d$  as follow:

- (i)  $AA'$  is a long axis of  $E$ ;
- (ii) all other semi-axes of  $E$  are equal of length  $h$ .

Indeed, consider the function of  $d$  variables:

$$F(x_1, \dots, x_d) := \frac{x_1^2}{h^2} + \dots + \frac{x_{d-1}^2}{h^2} + \frac{x_d^2}{R^2},$$

then  $F(x_1, \dots, x_d) = 1$  gives the hyperellipsoid when  $AA'$  is lying in the  $x_d$ -axis. Generally, if the line  $AA'$  has a unit directional vector  $\vec{u}_d$ , extend it to a orthonormal basis  $\beta := \{\vec{u}_1, \dots, \vec{u}_{d-1}, \vec{u}_d\}$  of  $\mathbb{R}^d$ . Then there exists a unitary transformation  $T : \mathbb{R}^d \rightarrow \mathbb{R}^d$  which sends  $\beta$  to the standard orthonormal basis

$\{(1, 0, \dots, 0), \dots, (0, \dots, 0, 1)\}$  such that  $T(\vec{u}_d) = (0, \dots, 0, 1)$ . Then the  $d-1$ -dimensional hyperellipsoid  $E$  has equation  $F(T(x_1, \dots, x_d)) = 1$ . See Figure 3.




FIGURE 3

Now let  $F \subset \mathbb{R}^d$  be the domain enclosed by  $E$  (including the points of  $E$ ). Apparently,  $F$  satisfies the condition (a) and (b) of Minkowski's Theorem. Moreover, it is known that the volume of the hyperellipsoid is

$$(3.2) \quad \frac{\pi^{\frac{d}{2}}}{\Gamma(\frac{d}{2} + 1)} h^{d-1} R,$$

here  $\Gamma$  is the gamma function, so

$$(3.3) \quad \Gamma(\frac{d}{2} + 1) = \frac{d}{2} \Gamma(\frac{d}{2}) = \frac{d}{2} (\frac{d}{2} - 1) \cdots \gamma_0,$$

where  $\gamma_0 = 1$  if  $d$  is even,  $\gamma_0 = \frac{\pi}{2}$  if  $d$  is odd. By Minkowski's Theorem, if we choose  $h$  such that

$$(3.4) \quad \frac{\pi^{\frac{d}{2}}}{\Gamma(\frac{d}{2} + 1)} h^{d-1} R = 2^d,$$

then  $F$  contains an integral point other than  $O$ . This implies that, if we set  $C = \frac{2^d \Gamma(\frac{d}{2} + 1)}{\pi^{\frac{d}{2}}}$ , and the tree radius  $r = CR^{\frac{1}{d-1}}$ , then any ray segment  $OA$  starting from  $O$  will be blocked by some tree at the integral point contained in  $F$  we constructed as above. Since  $\rho < r$ , that we complete the proof.  $\square$

Combining Theorem 1 and Theorem 2, we obtain the main result of this article:

**THEOREM 3.** *For  $d$ -dimensional orchard visibility problem, as the radius of orchard  $R$  goes to infinity,*

$$(3.5) \quad \rho = O(R^{-\frac{1}{d-1}}).$$

## 4. SOME FURTHER THOUGHTS

We can still ask a lot of questions concerning the orchard visibility problem in arbitrary dimension. For example, for  $d = 2$ , it has been proved in [2] that  $\rho = \frac{1}{R}$ , or

$$(4.1) \quad \lim_{R \rightarrow \infty} \rho R = 1.$$

Inspired by our results, it is natural to ask if we can find a constant  $l$  for dimension  $d$  such that

$$(4.2) \quad \lim_{R \rightarrow \infty} \rho^{d-1} R = l.$$

However, our estimation in this article using polyhedra is apparently not precise and fine enough for such a conclusion. We will explore this problem in the future.

## REFERENCES

- [1] Clyde P. Kruskal *The orchard visibility problem and some variants*, Journal of computer and system sciences, 74 (2008) 587–597.
- [2] Thomas Tracy Allen *Polya's Orchard Problem*, The American Mathematical Monthly, Vol. 93, No. 2 (1986) 98-104.
- [3] Alexandru Hening and Michael Kelly *On Polya's Orchard Problem*, Rose-Hulman Undergraduate Mathematics Journal: Vol. 7 : Iss. 2 , Article 9.
- [4] I. G. Macdonald *The volume of a lattice polyhedron*, Mathematical Proceedings of the Cambridge Philosophical Society, 59 (1963) 719-726.